

The Petersen graph is the smallest 3-cop-win graph

Andrew Beveridge*, Paolo Codenotti†, Aaron Maurer‡, John McCauley§, and Silviya Valeva¶

April 3, 2012

Abstract

In the game of *cops and robbers* on a graph $G = (V, E)$, k cops try to catch a robber. On the cop turn, each cop may move to a neighboring vertex or remain in place. On the robber's turn, he moves similarly. The cops win if there is some time at which a cop is at the same vertex as the robber. Otherwise, the robber wins. The minimum number of cops required to catch the robber is called the *cop number* of G , and is denoted $c(G)$. Let m_k be the minimum order of a connected graph satisfying $c(G) \geq k$. Recently, Baird and Bonato determined via computer search that $m_3 = 10$ and that this value is attained uniquely by the Petersen graph. Herein, we give a self-contained mathematical proof of this result. Along the way, we give some characterizations of graphs with $c(G) > 2$ and very high maximum degree.

1 Introduction

Let $G = (V, E)$ be a simple undirected graph on n vertices. The game of cops and robbers on G was independently introduced by Quilliot [18] and Nowakowski and Winkler [15]. The game is played between k cops $C_1, C_2, \dots, C_k$ and one robber R . First, the cops are placed at k vertices of the graph. Then the robber is placed on a vertex. During play, the cops and the robber move alternately. On the cop turn, each cop may move to a neighboring vertex or remain in place. The cops can coordinate their strategy and multiple cops may occupy the same vertex. On the robber's turn, he moves similarly. This is a full information game, in the sense that the locations of the cops and robber are always known to all players. The cops win if there is some finite time at which a cop is colocated with the robber. Otherwise, the robber wins. The minimum number of cops required to catch the robber (regardless of robber's strategy) is called the *cop number* of G , and is denoted $c(G)$. When $c(G) = k$, we say that G is *k-cop-win*.

*Department of Mathematics, Statistics and Computer Science, Macalester College, Saint Paul MN 55105

†Institute for Mathematics and its Applications, University of Minnesota, Minneapolis MN 55455

‡Department of Mathematics, Carleton College, Northfield MN 55057

§Department of Mathematics, Haverford College, Haverford PA 19041

¶Department of Mathematics, University of Iowa, Iowa City IA 52242

The game of cops and robbers has received considerable attention in recent years; for an introduction see the surveys [2, 11] and the monograph [6]. The most important open question is Meyniel's conjecture that for every graph G over n vertices, $c(G) = O(\sqrt{n})$. The history of Meyniel's conjecture is surveyed in [3], and the best known bound of $c(G) \leq n^{2^{-(1+o(1))\sqrt{\log n}}}$ was obtained independently in [12, 19, 10]. Various authors have recently studied the cop number for random graph models [4, 13, 16, 7, 5]. In particular, Prałat and Wormald [17] have shown that Meyniel's conjecture holds for the Erdős-Renyi random graph model.

The original papers [18, 15] characterized the graphs for which $c(G) = 1$. Given a vertex v , its *neighborhood* is $N(v) = \{u \in V \mid (v, u) \in E\}$, and its *closed neighborhood* is $\overline{N}(v) = \{v\} \cup N(v)$. A vertex v is *dominated* by the vertex w if $\overline{N}(v) \subseteq \overline{N}(w)$. A dominated vertex is also called a *pitfall* or *corner*. A graph G is *dismantleable* if we can reduce G to a single vertex by successively removing dominated vertices. These papers prove that a connected graph G has $c(G) = 1$ if and only if G is dismantleable.

Aigner and Fromme [1] introduced the cop number as described above. In addition to proving that if G is planar then $c(G) \leq 3$, they establish the following useful result. Let $\delta(G)$ denote the minimum degree of G . Recall that the *girth* $g(G)$ of a graph is the size of its smallest cycle (if G is acyclic then $g(G)$ is infinite). They proved that if G is a graph with finite girth $g(G) \geq 5$ then $c(G) \geq \delta(G)$. The Petersen graph H is a 10 vertex, 3-regular graph of girth 5. Aigner and Fromme's result guarantees that the $c(H) \geq 3$, and it shows a winning 3-cop strategy.

Let m_k denote the minimum order of a connected graph with $c(G) \geq k$. Clearly, $m_1 = 1$ and $m_2 = 4$. Recently, Baird and Bonato [3] used a computer search to prove that $m_3 = 10$ and that this value is attained uniquely by the Petersen graph. We give a self-contained mathematical proof of this result, split into two theorems.

Theorem 1.1. *If G is a connected graph on at most 9 vertices, then $c(G) \leq 2$.*

Theorem 1.2. *The Petersen graph is the unique connected graph on 10 vertices that requires 3 cops. All other connected graphs of order 10 are 2-cop-win.*

Our proofs follow from a series of observations and lemmas about the cop number of graphs with very large maximum degree. It is obvious that a graph with a *universal vertex* v such that $\deg(v) = n - 1$ is cop-win. We prove some structural results concerning graphs containing a vertex whose codegree is a small constant.

Lemma 1.3. *Let G be a connected graph on n vertices. If there is a node $u \in V(G)$ of degree at least $n - 6$, then either $c(G) \leq 2$ or the induced subgraph $G[V - \overline{N}(u)]$ is a 5-cycle.*

Corollary 1.4. *If $\Delta(G) \geq n - 5$ then $c(G) \leq 2$.*

Lemma 1.3 and its immediate corollary are crucial tools in proving our main results. In particular, Theorem 1.1 is a quick consequence of Corollary 1.4. This reduces our search to 10 vertex graphs with $2 \leq \delta(G) \leq \Delta(G) \leq 4$. It takes some additional effort to show that $c(G) = 3$ forces $\Delta(G) = 3$. At that point, we need the following lemma.

Lemma 1.5. *Let G be a graph with a vertex u with $\Delta(G) = \deg(u) = n - 7$ and such that $\deg(v) \leq 3$ for every $v \in V - \overline{N}(u)$. Then either $c(G) \leq 2$ or the induced subgraph $G[V - \overline{N}(u)]$ is a 6-cycle.*

This lemma can be generalized a bit more. In particular, if we remove the restriction on the maximum of degree of vertices in $V - \overline{N}(u)$, then the proofs of Lemmas 1.3 and 1.5 can be adapted to show that H must contain an induced 5-cycle or 6-cycle. However, the case analysis is cumbersome, so we have opted for this simpler formulation. The version stated above is sufficient to prove our main result: that the Petersen graph is the only 10-vertex graph requiring 3 cops.

We conclude this section with some reflections on our main results. The Petersen graph is the unique 3-regular graph of girth 5 of minimal size, so that Theorem 1.2 provides a tight lower bound for n when $c(G) = 3$. We wonder whether a similar result holds for general cop numbers, and we formulate some open question in this vein. Recall that a (k, g) -cage is a k -regular graph with girth g of minimal order. For a survey of cages, see [9]. The Petersen graph is the unique $(3, 5)$ -cage, and in general, cages exist for any pair $k \geq 2$ and $g \geq 3$.

As discussed earlier, Aigner and Fromme [1] proved that graphs with girth 5, and degree k have cop number at least k ; in particular, if G is a $(k, 5)$ -cage then $c(G) \geq k$. Let $n(k, g)$ denote the order of a (k, g) -cage. Is it true that a $(k, 5)$ -cage is k -cop-win? Next, since we have $m_k \geq n(k, 5)$, it is natural to wonder whether $m_k = n(k, 5)$ for $k \geq 4$. It seems reasonable to expect that this is true at least for small values of k . It is known that $n(4, 5) = 19$, $n(5, 5) = 30$, $n(6, 5) = 40$ and $n(7, 5) = 50$. Do any of these cages attain the analogous m_k ?

More generally, we can ask the same question for large k : is m_k achieved by a $(k, 5)$ -cage? If Meyniel's conjecture is true, then $m_k = \Omega(k^2)$. Meanwhile, it is known that $n(k, 5) = \Theta(k^2)$, so an affirmative resolution would be consistent with Meyniel's conjecture. We note that Baird and Bonato [3] have already observed that Meyniel's conjecture implies that $m_k = \Theta(k^2)$, using a projective plane construction to obtain the upper bound, rather than the existence of $(k, 5)$ -cages.

2 Preliminaries

2.1 Definitions and Notation

For a graph G , we will denote by $V(G)$ the vertex set of G , and $E(G)$ the edge set of G . If the graph G is clear from context, we will sometimes use V for $V(G)$ and E for $E(G)$. We also use the notation $v(G) := |V(G)|$. For $S \subseteq V$, the *induced subgraph* on S , denoted $G[S]$, is the graph with vertex set S , and edge set $E(G[S]) = \{(u, v) \in E(G) \mid u, v \in S\}$. For sets S, T , we denote the difference of T by S by $T - S := \{t \in T \mid t \notin S\}$.

We specify some additional vertex notation. For $u, v \in V$, we write $u \sim v$ when $(u, v) \in E$. For $S \subset V$, we write $u \sim S$ when $u \notin S$ and there exists $v \in S$ such that $u \sim v$. We define $N(S) = \cup_{v \in S} N(v) - S$ and $\overline{N}(S) = \cup_{v \in S} \overline{N}(v)$. For convenience, we set $N(u, v) = N(\{u, v\})$. For sets of vertices $S, T \subseteq V$, we denote the set of edges between the two sets by $[S : T] := \{(s, t) \in E \mid s \in S, t \in T\}$, and we use $|S : T|$ to denote the size of this edge set. We denote the *degree* of a vertex u by $\deg(u)$ and denote the minimum degree by $\delta(G) = \min_{v \in V(G)} \deg(v)$ and the maximum degree by $\Delta(G) = \max_{v \in V(G)} \deg(v)$. We generalize the latter symbol to subsets of vertices: for $S \subseteq V$, $\Delta(S) = \max_{s \in S} \deg(s)$.

We now introduce our notation for the state of the game. We fix a connected graph G on which the game is played. The state of the game is a pair $(C; r)$, where G is a connected graph, C is a k -tuple of vertices $C = (c_1, c_2, \dots, c_k)$, where $c_i \in V(G)$ is the current position of cop C_i , and $r \in V(G)$ is the current position of the robber R . For notational convenience, we write $(c_1, \dots, c_k; r)$ for $((c_1, \dots, c_k); r)$. When we need to specify whose turn it is to act, we underline the position of the player whose turn it is: $(\underline{C}; r)$ denotes that it is the cops' turn to move, and $(C; \underline{r})$ the robber's.

We use a shorthand notation to describe moves: $(\underline{c_1, \dots, c_k}; r) \rightarrow (\underline{c'_1, \dots, c'_k}; \underline{r})$ denotes the cop move where each C_i moves from c_i to c'_i . Similarly $(c_1, \dots, c_k; \underline{r}) \rightarrow (\underline{c_1, \dots, c_k}; r')$ denotes the robber's move from r to r' . We will concatenate moves and we use the shorthand $\rightarrow$, meaning a cop move followed by a robber move:

$$(\underline{c_1, \dots, c_k}; r) \rightarrow (\underline{c'_1, \dots, c'_k}; r') \equiv (\underline{c_1, \dots, c_k}; r) \rightarrow (\underline{c'_1, \dots, c'_k}; \underline{r}) \rightarrow (\underline{c'_1, \dots, c'_k}; r').$$

There will be cases where the strategy allows for either the robber or the cops to be in one of several positions. In general, for $T_i \subseteq V$, $S \subseteq V$, the state of the game has the form $(T_1, \dots, T_k; S)$ means that $c_i \in T_i$, and $r \in S$.

The robber's *safe neighborhood*, denoted $S(R)$, is the connected component of $G - \overline{N}(C)$ con-

taining the robber. We say that the robber is *trapped* when $S(R) = \emptyset$. This condition is equivalent to having both $r \in N(C)$ and $N(r) \subset \overline{N}(C)$. Once the robber is trapped, he will be caught on the subsequent cop move, regardless of the robber's next action. When the robber is trapped, we are in a *cop-winning position*, denoted by $\mathcal{C}$.

2.2 The end game

We frequently use the following facts to identify cop-win strategies for two cops in the end game. We state a more general version of these results for k cops.

We need the following property of a cop-win graphs, which first appears in [8]. Every cop-win graph has at least one cop-winning *no-backtrack strategy*, meaning that the cop never repeats a vertex during the pursuit. Typically, a graph has multiple no-backtrack strategies. We say that a vertex v is *no-backtrack-winning* if there is a cop-winning no-backtrack strategy starting at v . For example, when G is a tree, every vertex is no-backtrack-winning.

Next we fix some notation. For a fixed set $U = \{u_1, \dots, u_t\} \subset V$, let $N'_U(u_j) := N(u_j) - \overline{N}(U - u_j)$ be the neighbors of u_j that are not adjacent to any other vertex in U .

Observation 2.1. *Let $(\underline{C}; r)$ be the state of the game. Suppose that there exists a $c_j \in C$ such that either (a) $[S(R) : N'_C(c_j)] = \emptyset$ and $G[S(R)]$ is cop-win; or (b) $N(S(R)) \cap N'_C(c_j) = \{v\}$ such that $H = G[S(R) + v]$ is cop-win and v is no-backtrack-winning in H ; then the cops can win from this configuration.*

Proof. Let $S = S(R)$ be the initial safe neighborhood of R . In both cases, only cop C_j is active, while the others remain stationary. In case (a), cop C_j moves into S and follows a cop-win strategy on $G[S]$. In case (b), cop C_j moves to v and then follows a no-backtrack strategy on $G[S + v]$. This prevents the robber from ever reaching v . In both cases, the only way for the robber to avoid capture by C_j is to step into the neighborhood of the remaining cops. $\square$

We highlight two useful consequences that are used heavily for $k = 2$ in our subsequent proofs.

Corollary 2.2. *Let $(\underline{C}; r)$ be the state of the game, played with $k \geq 2$ cops. If $|S(R)| \leq 2$ and $|N(S(R))| \leq 2k - 1$, then the cops can win.*

Proof. Let $S = S(R)$. We have $|N(S) \cap N(C)| \leq 2k - 1$, so the pigeon hole principle ensures that there exists a cop C_j such that $|N(S) \cap N'_C(c_j)| \leq 1$. We are done by Observation 2.1, since every

vertex of a connected 2-vertex graph is non-backtrack-winning. $\square$

Corollary 2.3. *Let $(\underline{C}; r)$ be the state of the game, played with $k \geq 2$ cops. If $\max_{v \in S(R)} \deg_G(v) \leq 3$ and $S(R)$ contains at most one vertex of degree 3, then the cops can win.*

Proof. Let $S = S(R)$. Since $G[S]$ is connected, we have $[S : N(C)] \leq 3$. Therefore, some cop C_j has $|S : N'_C(c_j)| \leq 1$. If $G[S]$ is a tree, then we are done by Observation 2.1. If $G[S]$ is not a tree, then $G[S]$ must be unicyclic with one degree 3 vertex, say u . Therefore, $|S : N(C)| = 1$, and except for u , every vertex in the cycle has degree 2 in G . A cop-winning strategy is as follows: two cops move until they both reach u . Now $S(R)$ is a path, so Observation 2.1 completes the proof. $\square$

3 Graphs with $\Delta(G) \geq n - 6$

In this section, we prove Lemma 1.3 and Theorem 1.1. We also make progress on the proof of Theorem 1.2 by showing that if $v(G) = 10$ and $\Delta(G) = 4$, then $c(G) \leq 2$. For convenience, we recall the statements of these results prior to their respective proofs.

(Lemma 1.3). *Let G be a connected graph on n vertices. If there is a node $u \in V(G)$ of degree at least $n - 6$, then either $c(G) \leq 2$ or the induced subgraph $G[V - \overline{N}(u)]$ is a 5-cycle.*

Proof. Let $H = G[V - \overline{N}(u)]$. By Observation 2.1(a), if H is cop-win, then $c(G) \leq 2$. In particular this holds if H does not contain an induced cycle of length at least 4. So we only need to consider the case where $v(H) = 4$ or 5, and H contains an induced 4-cycle. Let x_1, x_2, x_3, x_4 form the 4-cycle in H (in that order). Let x_5 be the additional vertex (if present).

We now distinguish some cases based on $N(x_5) \cap H$. If $x_5 \sim x_i$ for every $i \in \{1, 2, 3, 4\}$, then H is cop-win, and hence $c(G) \leq 2$. This leaves us with 5 cases to consider, depicted in Figure 3.1. Case (a) includes the situation when $\deg(u) = n - 5$, and there is no vertex x_5 .




Figure 3.1: The five cases for $G[V - \overline{N}(u)]$ in Lemma 1.3.




Figure 3.2: The four classes of possible structures of G for Claim 3.2. Vertex x_5 might not be present, and dashed edges might not be present.

First we make some technical observations. We start by noting that moving to x_5 is in most situations a bad idea for the robber in Cases (a), (b) and (c).

Claim 3.1. *In Cases (a), (b), and (c), if the state of the game is of the form $(\overline{N}(u), V(H); x_5)$, the cops have a winning strategy.*

Proof. C_1 moves to u . In Case (a), $S(R) = \{x_5\}$ and we are already done by Corollary 2.2. In Case (b), if possible, C_2 moves directly to x_2 ; otherwise C_2 moves first to x_1 and then to x_2 ; in either case the robber is trapped at x_5 . In Case (c), C_2 moves to x_2 or x_1 (whichever c_2 is adjacent to), again trapping the robber in x_5 . $\square$

Next we look at the structure of $N(y) \cap V(H)$ for nodes $y \in N(u)$.

Claim 3.2. *Suppose the state of the game has the form $(\overline{N}(u), \{x_1, x_3\}; y)$, where $y \in N(u)$ is such that either (a) $N(y) \cap V(H) = \{x_2, x_4\}$, or (b) y is adjacent to at most one of x_2 or x_4 ; then the cops have a winning strategy.*

Proof. Figure 3.2 shows the four classes of possible graph structures. Let us first consider the structure (B1). Let $z = x_2$. C_1 moves to u , and C_2 moves to z . Now the robber is trapped in all cases of Figure 3.1 except Case (a). In Case (a) the robber's only move is to x_5 . After this move, the robber can be caught by Claim 3.1. The same cop strategy works for structures (B2) and (B3), taking $z = x_4$. A simplified version of this proof shows that the same cop strategy works for structure (A), taking $z = c_2$. $\square$

REMARK: in Cases (d) and (e) of Figure 3.1, x_5 and x_1 are symmetric, so the statement holds also for configuration $(\overline{N}(u), x_5; y)$

Our next observation concerns the situation where there are two nodes in $N(u)$ that do not satisfy the condition of the previous claim.

Claim 3.3. *If there are two vertices $y, z \in N(u)$ such that $\{x_2, x_3, x_4\} \subset N(y)$, and $\{x_1, x_2, x_4\} \subset N(z)$, then $c(G) \leq 2$.*

Proof. First we deal with all cases but Case (d). The cops start at u and z . If the robber starts at x_3 , the cops' winning strategy is: $(\underline{u}, z; x_3) \rightarrow (u, y; \underline{x_3}) \mathcal{C}$. If the robber starts at x_5 , the strategy will depend on the structure of H . In Cases (a),(b),(c), we are done by Claim 3.1. In Case (e), the following is a winning strategy: $(\underline{u}, z; x_5) \rightarrow (u, x_1; \underline{x_5}) \mathcal{C}$.

The remainder of the proof deals with Case (d), which requires a more involved argument.

First suppose that there exists $w \in N(u)$ such that $\{x_2, x_4, x_5\} \subset N(w)$. Then the cops start at u and z . The robber can start at x_3 or x_5 in either case the cops have a winning strategy: $(\underline{u}, z; x_3) \rightarrow (y, u; \underline{x_3}) \mathcal{C}$; or $(\underline{u}, z; x_5) \rightarrow (w, u; \underline{x_5}) \mathcal{C}$.

Now assume that no such w exists. Start the cops at u and y . The robber starts in $\{x_1, x_5\}$. If the robber starts at x_1 , then $(\underline{u}, y; x_1) \rightarrow (y, u; \underline{x_1}) \mathcal{C}$. So we may assume the robber starts at x_5 . If $|N(x_5) \cap N(u)| \leq 1$, we are done by Corollary 2.2. Otherwise, the cops move by $(\underline{u}, x_3; x_5) \rightarrow (v, z; \underline{x_5})$, for some $v \in N(x_5) \cap N(u)$. The robber is forced to move to some $w \in N(x_5) \cap N(u)$ (if no such w exists, then R is trapped). By our initial argument, w cannot be adjacent to both x_2 and x_4 , but then the state satisfies the conditions of Claim 3.2(b), and we are done. $\square$

Claim 3.4. *Either $c(G) \leq 2$, or we can relabel the vertices of H via an automorphism of H so that x_1 is adjacent to $N(u)$.*

Proof. Suppose that no such relabeling exists. We will show a winning strategy for the cops, starting at u and x_3 . In Cases (a) and (b) the claim follows from Corollary 2.2 (either $S(R) = \{x_1\}$ or $S(R) = \{x_5\}$). In Cases (d) and (e), $S(R) \subseteq \{x_1, x_5\}$, and we are assuming that both x_1 and x_5 have no edges to $N(u)$; hence $|N(S(R))| \leq 2$, and we are again done by Corollary 2.2. In Case (c) $S(R) = \{x_1, x_5\}$, and we are assuming that both x_1 and x_2 do not have neighbors in $N(u)$. By Claim 3.1, we may assume R does not start at x_5 , and so R starts at x_1 . Let $v \in N(u) \cap N(x_5)$ (if

$x_5 \approx N(u)$, then $N(S(R))$ is dominated by $c_2 = x_3$). Now the cops can win by following the strategy: $(\underline{u}, x_3; x_1) \rightarrow (\underline{v}, x_3; x_1) \notin \mathcal{C}$. $\square$

Armed with the above claims, we now conclude the proof Lemma 1.3.

By Claim 3.4, we may assume $x_1 \sim w \in N(u)$. Initially place C_1 at u and C_2 at x_1 . The robber could start at x_3 or, in Cases (a), (b), and (d), at x_5 . If the robber starts at x_5 in Cases (a) and (b), then we are done by Claim 3.1. In Case (d), x_5 and x_3 are symmetric, so without loss of generality, $r = x_3$, and the initial state is $(\underline{u}, x_1; x_3)$.

If $x_3 \approx N(u)$, then the cops win by Corollary 2.2. Otherwise let $v \in N(x_3) \cap N(u)$. Then C_1 moves from u to v , while C_2 remains fixed at x_1 , forcing R to some $y \in N(u) \cap N(x_3)$, with $y \approx v, y \approx x_1$. If no such y exists, then R is trapped. If y is adjacent to only one of x_2 or x_4 , we are in the state $(\underline{v \in N(u)}, x_1; y)$, which satisfies the conditions of Claim 3.2(b), and hence the cops have a winning strategy.

Otherwise y is adjacent to x_2, x_3 , and x_4 . The cops move $(\underline{v}, x_1; y) \rightarrow (x_3, w; \underline{y})$, for some $w \in N(x_1) \cap N(u)$. If $y \sim x_5$, and R moves to x_5 , then the cops win: in Cases (a),(b),(c) we are done by Claim 3.1; in Case (d), $(x_3, w; x_5) \rightarrow (y, u; \underline{x_5}) \notin \mathcal{C}$; in Case (e), the cops can adopt a different strategy from the beginning: $(\underline{u}, y; x_1) \rightarrow (u, x_5; \underline{x_1}) \notin \mathcal{C}$. The only other option is for R to move to some $z \in N(u), z \approx x_3$. So the state is $(x_3, w; z)$. Either the pair y, z satisfies the conditions of Claim 3.3, or the current state satisfies the conditions of Claim 3.2(b) or (a). In either case, we are done. This concludes the proof of Lemma 1.3. $\square$

We now state some quick but useful consequences of Lemma 1.3.

Corollary 3.5. *Let G be a graph on n vertices. If there is a vertex $u \in V$ of degree at least $n - 6$, and a vertex $v \in V - \overline{N}(u)$ such that $|N(v) - N(u)| \geq 3$, then $c(G) \leq 2$.*

Proof. v has three neighbors in $G[V - \overline{N}(u)]$, and hence $G[V - \overline{N}(u)]$ cannot be a 5-cycle. $\square$

Corollary 3.6. *Let G be a graph on n vertices. If there is a vertex u of degree at least $n - 6$ and a vertex $v \in V - \overline{N}(u)$ with $\deg(v) \leq 3$, then $c(G) \leq 2$.*

Proof. By Lemma 1.3, we only need to consider the case where $G[V - \overline{N}(u)]$ is a 5-cycle, x_1, x_2, x_3, x_4, x_5 (in that order). Without loss of generality, let $\deg(x_1) \leq 3$, and $\deg(x_2) \geq 3$. For each $i = 1, \dots, 5$ such that $\deg(x_i) \geq 3$, pick some $y_i \in N(x_i) \cap N(u)$ arbitrarily (we allow

$y_i = y_j$ for $i \neq j$). The game starts as $(\underline{u}, \underline{x_4}; \{x_1, x_2\}) \rightarrow (\underline{u}, \{x_3, x_4\}; x_1)$. First we deal with the case where $\deg(x_1) = 2$ and the case where $\deg(x_1) = 3$ and $y_1 \sim x_4$. The cops' winning strategy for these two cases is the same: $(\underline{u}, \{x_3, x_4\}; x_1) \rightarrow (\underline{y_2}, \underline{x_4}; x_1) \rightarrow (x_2, x_4; \underline{x_1}) \mathcal{C}$.

Now we may assume that all x_i have degree 3, and hence y_i exists for all i . We may further assume that $x_4 \neq y_1$, and, since x_3 and x_4 are symmetric, we are also done in the case $y_1 \sim x_3$. The only remaining possibility is $N(y_1) \cap (V - \overline{N}(u)) \subseteq \overline{N}(x_1)$. Since x_3 and x_4 are symmetric, without loss of generality, the state is $(\underline{u}, \underline{x_4}; x_1)$. The cops first move to y_2 and x_5 , forcing the robber to y_1 , then in one more move, the robber is trapped at y_1 : $(\underline{y_2}, \underline{x_5}; y_1) \rightarrow (\underline{u}, x_1; \underline{y_1}) \mathcal{C}$. $\square$

These corollaries are enough to prove that every connected 9-vertex graphs is 2 cop-win, and to show that if $v(G) = 10$ and $\Delta(G) = 4$ then $c(G) \leq 2$.

(Theorem 1.1). *If G is a connected graph on at most 9 vertices, then $c(G) \leq 2$.*

Proof. If $\Delta(G) \geq 4$, then we are done by Lemma 1.3. If $\Delta(G) = 3$, then we are done by Corollary 3.6. $\square$

Lemma 3.7. *If $v(G) = 10$ and $\Delta(G) \geq 4$, then $c(G) \leq 2$.*

Proof. Let $u \in V(G)$ have degree at least 4. By Lemma 1.3, either $c(G) \leq 2$ or $\deg(u) = 4$, and $G[V - \overline{N}(u)]$ is a 5-cycle. Now, by Corollary 3.6, either $c(G) \leq 2$, or every $u \in V - \overline{N}(u)$ has $\deg(u) \geq 4$. In the latter case, $|[N(u) : V - \overline{N}(u)]| \geq 10$, thus, by the pigeon hole principle, there exists $v \in N(u)$ such that $|N(v) \cap (V - \overline{N}(u))| \geq 3$. We now deal with this case, namely u and v have degree 4, and $N(u) \cap N(v) = \emptyset$.

By Lemma 1.3, both $G[V(G) - \overline{N}(u)]$ and $G[V(G) - \overline{N}(v)]$ are 5-cycles. The resulting graph structure must be one of the two shown in Figure 3.3. Considering the structure in Figure 3.3(a), we note that $\deg(z_1) = \deg(z_2) = 3$ in order to maintain the induced 5-cycle structures, and hence we are done by Corollary 3.6.

Now suppose that G has the structure in Figure 3.3(b). In this case we show $\deg(x_3) = 3$, and we are again done by Corollary 3.6. To show that $\deg(x_3) = 3$, we look at each potential additional edge, and show that $V - \overline{N}(x_3)$ is not a 5-cycle, and hence we are done by Lemma 1.3. We only need to consider edges to y_1, y_2 or y_3 : other potential edges would not maintain the induced 5-cycle structure. We have $x_3 \sim y_1$ because $\{v, y_2, y_3\}$ form a triangle. We have $x_3 \sim y_3$ because z_1 is adjacent to

each of x_1, y_1, y_2 . Finally, $x_3 \sim y_2$ because the existence of this edge would force $y_3 \sim x_1$, which is symmetric to the forbidden $x_3 \sim y_1$. $\square$




Figure 3.3: The two possible starting structures in the proof of Lemma 3.7. Circled vertices cannot have additional edges.

4 Graphs with $\Delta(G) = n - 7$

In this section, we prove Lemma 1.5 and complete the proof of Theorem 1.2.

(Lemma 1.5). *Let G be a graph with a vertex u with $\Delta(G) = \deg(u) = n - 7$ and such that $\deg(v) \leq 3$ for every $v \in V - \overline{N}(u)$. Then either $c(G) = 2$ or the induced subgraph $G[V - \overline{N}(u)]$ is a 6-cycle.*

Proof. Let $H = G[V - \overline{N}(u)]$ and suppose that $c(G) > 2$. First, we observe that H must be connected. Otherwise, we can adapt the proof of Corollary 3.6 to show that $c(G) = 2$. Indeed, H has at most one component H_1 whose cop number is 2. We use the strategy described in the proof of Corollary 3.6 to capture the robber. The only alteration of the strategy is to address the robber moving from $N(u)$ to $H - H_1$. However, $|V(H - H_1)| \leq 2$, so this component is cop-win. One cop responds by moving to u , while the other moves into $H - H_1$ for the win (Observation 2.1(a)).

Therefore, we may assume that H is connected and $c(H) \geq 2$. This means that H must contain an induced k -cycle for $k \in 4, 5, 6$. Suppose that G contains an induced 4-cycle x_1, x_2, x_3, x_4 . Without loss of generality, $x_5 \sim x_1$, and x_6 is adjacent to at most three of $\{x_2, x_3, x_4, x_5\}$ (because we already have $\deg(x_1) = 3$). Start the cops at u and x_1 , so that $S(R)$ is one of $\{x_3\}$, $\{x_6\}$ or $\{x_3, x_6\}$. In the first two cases, $\Delta(S(R)) \leq 3$ so the cops win by Corollary 2.3. The last option occurs when $x_3 \sim x_6$. If x_6 has at most one neighbor in $N(u)$, we are again done by Corollary 2.3, since $\Delta(S(R)) \leq 3$. When x_6 has two neighbors in $N(u)$, the game play depends on the initial location of the robber. If the

robber starts at x_6 then C_1 holds at u while C_2 moves from x_1 to x_2 to x_3 , trapping the robber. If the robber starts at x_3 , then the roles are reversed: C_1 moves to x_6 in two steps while C_2 holds at x_1 . At this point, the robber is trapped.

Next, suppose that G contains an induced 5-cycle x_1, x_2, x_3, x_4, x_5 . Without loss of generality, $x_6 \sim x_1$. If x_6 is adjacent to two of the x_i , then we can place C_1 at u and C_2 at some x_j so that $|N(S(R)) \cap N(u)| \leq 1$, giving a cop winning position by Observation 2.1(b). Indeed, by symmetry there are only 2 cases to consider: if $x_6 \sim x_2$, then C_2 starts at x_4 and $S(R) = \{x_1, x_2, x_6\}$; if $x_6 \sim x_3$, then C_2 starts x_3 , and $S(R) = \{x_1, x_5\}$. So we may assume that x_6 has no additional neighbors in H . There are two cases to consider. If x_2 and x_4 do not share a neighbor in $N(u)$, then our game play begins with C_2 chasing R onto x_2 : $(u, x_1; \{x_3, x_4\}) \rightarrow \cdots \rightarrow (u, \{x_4, x_5\}; x_2)$. If x_2 is not adjacent to $N(u)$, then the cops can ensure $S(R)$ satisfies Corollary 2.3 on their next move. Indeed, C_2 moves to x_4 . If $N(x_6) \cap N(u) = \emptyset$, then the situation already satisfies Corollary 2.3, otherwise C_1 moves to $N(x_6) \cap N(u)$, and now the situation satisfies Corollary 2.3.

The final case to consider is when x_2 and x_4 are both adjacent to $y \in N(u)$. By symmetry, x_3 and x_5 are adjacent to $z \in N(u)$. By symmetry, there is one game to consider: $(u, x_1; x_3) \rightarrow (z, x_2; x_4)$ which is cop-win by Corollary 2.3. Thus, the only option for H is an induced 6 cycle. $\square$

We can now prove that the Petersen graph is the unique 3 cop-win graph of order 10.

Observation 4.1. *The Petersen graph is the only 3-regular graph G such that for every vertex $u \in V(G)$, $G[V(G) - \overline{N}(u)]$ is a 6-cycle.*

Proof. This is easily checked against the 18 possible 3-regular graphs of order 10 listed at [14], but here we give a direct proof. Pick any vertex u . The complement is a 6-cycle, where every vertex is adjacent to exactly one vertex in $N(u)$. Let $N(u) = \{y, z, w\}$. Now pick a vertex x_1 on the 6-cycle, $x_1 \sim y$. Because $V - N(x_1)$ must form a 6-cycle, we must have that $x_3 \sim w$ and $x_5 \sim z$ (by symmetry this is the only option). The only remaining edges to add are a matching between x_2, x_4, x_6 and y, z, w . To avoid a triangle in $V - N(y)$, we cannot have $x_4 \sim y$ or $x_4 \sim z$, hence $x_4 \sim w$, and, by symmetry, $x_2 \sim z$, and $x_6 \sim w$. This is the Petersen graph. $\square$

(Theorem 1.2). *The Petersen graph is the unique connected graph on 10 vertices that requires 3 cops. All other connected graphs of order 10 are 2-cop-win.*

Proof. Let G be a connected graph of order 10 such that $c(G) = 3$. We have $\delta(G) \geq 2$: otherwise the

leaf $v \in V(G)$ is a dominated vertex, so $c(G) = c(G - v) \leq 2$ by Theorem 1.1. Lemma 3.7 ensures that $\Delta(G) \leq 3$. Clearly $\Delta(G) = 3$ since a connected 2-regular graph is a cycle which is 2-cop-win.

Suppose a vertex $u \in V(G)$ has $\deg(u) = 3$. Then, by Lemma 1.5, $G[V - \overline{N}(u)]$ must be a 6-cycle. If every vertex in $N(u)$ has degree 3, then G is 3-regular with $c(G) = 3$, and therefore G is the Petersen graph by Fact 4.1. Otherwise there is a vertex $x_1 \in V - \overline{N}(u)$ with $\deg(x_1) = 2$. In the rest of the proof we give a winning strategy for the cops in this case.

Let the 6-cycle $G[V - \overline{N}(u)]$ be $\{x_1, x_2, x_3, x_4, x_5, x_6\}$. Without loss of generality, $\deg(x_1) = 2$ and $\deg(x_2) = 3$. Let $k = \max\{i \mid \deg(x_i) = 3\}$. The initial configuration is $(\underline{u}, x_4; \{x_1, x_2, x_6\})$. If $k \leq 5$ then the cops win by Corollary 2.3. When $k = 6$, our strategy depends on the initial robber location. Let $y \in N(u) \cap N(x_2)$. We either have $(\underline{u}, x_4; x_2) \rightarrow (y, x_4; x_1) \rightarrow (y, x_5; x_1) \mathcal{C}$, or $(\underline{u}, x_4; x_1) \rightarrow (y, x_5; x_1) \mathcal{C}$, or $(\underline{u}, x_4; x_6) \rightarrow (y, x_5; x_1) \rightarrow (y, x_6; x_1) \mathcal{C}$. The robber is trapped for every initial placement. $\square$

5 Acknowledgments

This work was undertaken at the Institute for Mathematics and its Applications: first as part of its Summer 2010 Special Program on Interdisciplinary Research for Undergraduates, and subsequently while the first two authors participated in the IMA 2011-2012 Special Year on the Mathematics of Information. We are grateful to the IMA for its generous support. We also thank Volkan Isler and Vishal Saraswat for helpful conversations.

References

- [1] M. Aigner and M. Fromme. A game of cops and robbers. *Discrete Appl. Math.*, 8:1–12, 1983.
- [2] B. Alspach. Sweeping and searching in graphs: a brief survey. *Mathematiche*, 59:5–37, 2006.
- [3] W. Baird and A. Bonato. Meyniel’s conjecture on the cop number: a survey. Submitted.
- [4] B. Bollobas, G. Kun, and I. Leader. Cops and robbers in random graphs. Submitted.
- [5] A. Bonato, G. Kemkes, and P. Prałat. Almost all cop-win graphs contain a universal vertex. Submitted.

- [6] A. Bonato and R. Nowakowski. *The Game of Cops and Robbers on Graphs*. American Mathematical Society, 2011.
- [7] A. Bonato, P. Prałat, and C. Wang. Pursuit-evasion in models of complex networks. *Internet Mathematics*, 4(4):419–436, 2007.
- [8] N. Clarke and R. Nowakowski. Cops, robber and traps. *Utilitas Mathematica*, 60:91–98, 2001.
- [9] G. Exoo and R. Jajcay. Dynamic cage survey. *Electronic Journal of Combinatorics*, Dynamic Survey DS16, May 2011. <http://www.combinatorics.org>.
- [10] A. Frieze, M. Krivelevich, and P.-S. Lo. Variations on cops and robbers. *J. Graph Theory*. To appear.
- [11] G. Hahn. Cops, robbers and graphs. *Tatra Mountain Mathematical Publications*, 36:163–176, 2007.
- [12] L. Lu and X. Peng. On Meyniel’s conjecture of the cop number. Submitted.
- [13] T. Łuczak and P. Prałat. Chasing robbers on random gaphs: zigzag theorem. *Random Structures and Algorithms*, 37:516–524, 2010.
- [14] B. McKay. Combinatorial data. <http://cs.anu.edu.au/people/bdm/data>, Dec. 2011.
- [15] R. Nowakowski and P. Winkler. Vertex-to-vertex pursuit in a graph. *Discrete Mathematics*, 43(2–3):235 – 239, 1983.
- [16] P. Prałat. When does a random graph have constant cop number? *Australasian Journal of Combinatorics*, 46:285–296, 2010.
- [17] P. Prałat and N. Wormald. Meyniel’s conjecture holds in random graphs. Submitted.
- [18] A. Quilliot. *Jeux et pointes fixes sur les graphes*. PhD thesis, Université de Paris VI, 1978.
- [19] A. Scott and B. Sudakov. A bound for the cops and robbers problem. *SIAM J. Discrete Math*. To appear.